



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ANNALS OF MATHEMATICS.

VOL. I.

JULY, 1884.

NO. 3.

CONVERGENCE-SUBTRAHENDS FOR THE TRIGONOMETRICAL FUNCTIONS EXPRESSED IN INFINITE SUMS.

By PROF. R. D. BOHANNAN, University of Virginia.

If $F(z)$ is a function of a complex variable, $z = x + iy$, in general finite and continuous, but infinite on the points, $z = a_1, a_2, a_3, \dots, a_n$, and so infinite that $\lim (z - a_n) F(z) = A_n$ when $z = a_n$, where A_n is not infinite, then

$$F(z) = \sum_1^n \frac{A_n}{z - a_n} + G(z),$$

where $G(z)$ is a function not infinite for any finite value of z . When the number of points on which $F(z)$ is infinite after the manner stated, is infinite, then

$$F(z) = \lim \sum_1^\infty \left(\frac{A_n}{z - a_n} + S(z) \right) + G'(z).*$$

For any particular function, that which offers the greatest difficulty is the determination of the functions $G(z)$ and $G'(z)$. For this there seems to be no general method, except in so far as the contour-integration method of Cauchy is applicable; and in the application of this, great care is necessary.†

In all cases where an infinite sum is present, $S(z)$, called, conveniently, the Convergence-Subtrahend, must be so determined as to render the infinite sum convergent. Generally a number of functions may be found which will do this, thus making $G'(z)$ also variable. For a new function, after $S(z)$ has been determined, $G'(z)$ is quite arbitrary; for an old function whose properties are well known, $G'(z)$ must be so determined, in connection with the infinite sum, as to bring out these properties.

*See Weierstrass's *Zur Theorie der eindeutigen analytischen Functionen*, Berlin, 1876; the writings of Mittag-Leffler based on this pamphlet; and Schering's *Das Anschliessen einer Function an algebraische Functionen in unendlich vielen Stellen*, Goettingen, 1880.

†Compare Hermite's *Cours* 1881-82, XIIe Leçon, and his note on same in *American Journal of Mathematics* for September, 1883.

The object of this paper is to give a convenient form to $S(z)$ for the functions

$$\frac{1}{\sin z}, \frac{1}{\cos z}, \tan z, \cot z.$$

The only *essentially* singular point which these functions have, is "the point at infinity." Consequently here the infinite sums must be made *absolutely* convergent for all finite values of absolute z , other than those for which the functions become infinite. It will be assumed in what follows, that, for each of these functions, $G'(z)$, in connection with the $S(z)$ which we shall find, is zero. This can be proven, either by contour-integration,* or by showing, on the assumption that $G'(z)=0$, that the absolutely convergent infinite sums have all the properties of the functions they are made to represent, and agree with them in value, on points selected at random.

$$\frac{1}{\sin z}.$$

$\frac{1}{\sin z} = \infty$ when $z = m\pi$, m being 0 or any positive or negative integer, and

$\lim \frac{z - m\pi}{\sin z} = (-1)^m$ when $z = m\pi$; so that

$$\frac{1}{\sin z} = \frac{1}{z} + \sum_1^{\infty} \left(\frac{(-1)^m}{z - m\pi} + S_1(z) \right) + \sum_1^{\infty} \left(\frac{(-1)^m}{z + m\pi} + S_2(z) \right).$$

The sums $\sum_1^{\infty} \frac{(-1)^m}{z - m\pi}$ and $\sum_1^{\infty} \frac{(-1)^m}{z + m\pi}$ are not absolutely convergent. The infinite sums in the value of $\frac{1}{\sin z}$ are most conveniently rendered absolutely convergent if we determine $S_1(z)$ and $S_2(z)$ so that

$$\frac{1}{\sin z} = \frac{1}{z} + \sum_1^{\infty} (-1)^m \left\{ \frac{1}{z - m\pi} + \frac{1}{m\pi} \right\} + \sum_1^{\infty} (-1)^m \left\{ \frac{1}{z + m\pi} - \frac{1}{m\pi} \right\}.$$

This is identical in value with the usual expression

$$\frac{1}{\sin z} = \frac{1}{z} + 2z \sum_1^{\infty} \frac{(-1)^m}{z^2 - m^2\pi^2},$$

but the former shows, almost on its face, the periodicity of $\sin z$; and the periodicity of the trigonometrical functions is their most interesting property.

*See Briot et Bouquet's *Fonctions Doublement Periodiques*, p. 123; also their *Fonctions Elliptiques*, p. 281; also Hermite's *Cours*, p. 79.

$$\frac{1}{\cos z}.$$

$$\frac{1}{\cos z} = \infty \text{ when } z = \pm (2m+1)\frac{\pi}{2}, \text{ and } \lim_{z \rightarrow \pm (2m+1)\frac{\pi}{2}} \frac{(2m+1)\frac{\pi}{2} - z}{\cos z} = (-1)^m.$$

Hence, corresponding to the formula for $\frac{1}{\sin z}$, we have

$$\begin{aligned} \frac{1}{\cos z} &= \frac{1}{\frac{\pi}{2} - z} + \sum_1^{\infty} (-1)^m \left(\frac{1}{\frac{\pi}{2} - z - m\pi} + \frac{1}{m\pi} \right) \\ &\quad + \sum_1^{\infty} (-1)^m \left(\frac{1}{\frac{\pi}{2} - z + m\pi} - \frac{1}{m\pi} \right). \end{aligned}$$

This formula shows the periodicity of $\cos z$ and the connection between $\sin z$ and $\cos z$.

Briot and Bouquet, both in their *Fonctions Doublement Periodiques*, p. 125, and in their *Fonctions Elliptiques*, p. 285, obtain, by grouping the terms of an infinite sum, the expression

$$\frac{1}{\cos z} = \pi \sum_0^{\infty} \frac{(2m+1)(-1)^m}{(2m+1)^2 \frac{\pi^2}{4} - z^2}.$$

This is not an allowable formula, for it is not absolutely convergent for any value of z , as may be readily shown.

Similar formulæ for $\tan z$ and $\cot z$ are

$$\tan z = \frac{1}{\frac{\pi}{2} - z} - \frac{1}{\frac{\pi}{2} + z} + \sum_1^{\infty} \left(\frac{1}{\frac{\pi}{2} - z + m\pi} - \frac{1}{m\pi} \right) + \sum_1^{\infty} \left(\frac{1}{\frac{\pi}{2} - z - m\pi} + \frac{1}{m\pi} \right)$$

and

$$\cot z = \frac{1}{z} + \sum_1^{\infty} \left(\frac{1}{z - m\pi} + \frac{1}{m\pi} \right) + \sum_1^{\infty} \left(\frac{1}{z + m\pi} - \frac{1}{m\pi} \right).$$